

- Colizzi, FS, **Beer, K**, Cuti, P, Deppisch, P, Martínez Torres, D, Yoshii, T, and Helfrich-Förster, C (2021). Antibodies against the clock proteins period and cryptochrome reveal the neuronal organization of the circadian clock in the pea aphid. *Front. Physiol.* *12*, 705048.
- Beer K**, Härtel S, Helfrich-Förster C (2021). The pigment-dispersing factor neuronal network systematically grows in developing honey bees. *J of Comparative Neurology*, 1-20.
- Beer K**, and Helfrich-Förster C (2020a). Post-embryonic development of the circadian clock seems to correlate with social life style in bees. *Front. Cell Dev. Biol.* *8*, 1-9.
- Beer K**, and Helfrich-Förster C (2020b). Model and non-model insects in chronobiology. *Front. Behav. Neurosci.* *14*, 1-23.
- Beer K**, Bloch G (2020). Circadian plasticity in honey bees. *The Biochemist*, *42*, 22–26.
- Menegazzi P, **Beer K**, Grebler V, Schlichting M, Schubert F K, and Helfrich-Förster C (2020). A functional clock within the main morning and evening neurons of *D. melanogaster* is not sufficient for wild-type locomotor activity under changing day length. *Front. Physiol.* *11*, 229.
- Beer K**, Schenk M, Helfrich-Förster C, and Holzschuh A (2019). The circadian clock uses different environmental time cues to synchronize emergence and locomotion of the solitary bee *Osmia bicornis*. *Sci. Rep.* *9*, 1, 17748.
- Beer K**, Kolbe E, Kahana NB, Yayon N, Weiss R, Menegazzi P, Bloch G and Helfrich-Förster C (2018) Pigment-dispersing factor-expressing neurons convey circadian information in the honey bee brain. *Open Biology*, *8*, 1, 170224.
- Beer K**, Joschinski J, Arrazola Sastre A, Krauss J, and Helfrich-Förster C (2017). A damping circadian clock drives weak oscillations in metabolism and locomotor activity of aphids (*Acyrtosiphon pisum*). *Sci. Rep.* *7*, 1, 14906.
- Fuchikawa T, **Beer K**, Linke-Winnebeck C, Ben-David R, Kotowoy A, Tsang VWK., Warman GR, Winnebeck EC, Helfrich-Förster C, and Bloch G (2017). Neuronal circadian clock protein oscillations are similar in behaviourally rhythmic forager honeybees and in arrhythmic nurses. *Open Biology*, *7*, 6, 170047.
- Joschinski J, **Beer K**, Helfrich-Förster C, and Krauss J (2016). Pea Aphids (Hemiptera: Aphididae) Have diurnal rhythms when raised independently of a host plant. *J. Insect Sci.* *16*, 31.
- Beer K**, Steffan-Dewenter I, Härtel S, and Helfrich-Förster C (2016). A new device for monitoring individual activity rhythms of honey bees reveals critical effects of the social environment on behavior. *J. Comp. Physiol. A* *202*, 555–565.